

This discussion paper is/has been under review for the journal Atmospheric Chemistry and Physics (ACP). Please refer to the corresponding final paper in ACP if available.

10 yr spatial and temporal trends of PM_{2.5} concentrations in the southeastern US estimated using high-resolution satellite data

X. Hu¹, L. A. Waller², A. Lyapustin³, Y. Wang^{3,4}, and Y. Liu¹

¹Department of Environmental Health, Rollins School of Public Health, Emory University, Atlanta, GA 30322, USA

²Department of Biostatistics & Bioinformatics, Rollins School of Public Health, Emory University, Atlanta, GA 30322, USA

³NASA Goddard Space Flight Center, Greenbelt, Maryland, USA

⁴University of Maryland Baltimore County, Baltimore, Maryland, USA

Received: 13 August 2013 – Accepted: 6 September 2013 – Published: 7 October 2013

Correspondence to: Y. Liu (yang.liu@emory.edu)

Published by Copernicus Publications on behalf of the European Geosciences Union.

ACPD

13, 25617–25648, 2013

10 yr trend of PM_{2.5}
concentrations in the
southeastern US

X. Hu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

Long-term $\text{PM}_{2.5}$ exposure has been reported to be associated with various adverse health outcomes. However, most ground monitors are located in urban areas, leading to a potentially biased representation of the true regional $\text{PM}_{2.5}$ levels. To facilitate epidemiological studies, accurate estimates of spatiotemporally continuous distribution of $\text{PM}_{2.5}$ concentrations are essential. Satellite-retrieved aerosol optical depth (AOD) has been widely used for $\text{PM}_{2.5}$ concentration estimation due to its comprehensive spatial coverage. Nevertheless, an inherent disadvantage of current AOD products is their coarse spatial resolutions. For instance, the spatial resolutions of the Moderate Resolution Imaging Spectroradiometer (MODIS) and the Multiangle Imaging Spectro-Radiometer (MISR) are 10 km and 17.6 km, respectively. In this paper, a new AOD product with 1 km spatial resolution retrieved by the multi-angle implementation of atmospheric correction (MAIAC) algorithm was used. A two-stage model was developed to account for both spatial and temporal variability in the $\text{PM}_{2.5}$ -AOD relationship by incorporating the MAIAC AOD, meteorological fields, and land use variables as predictors. Our study area is in the southeastern US, centered at the Atlanta Metro area, and data from 2001 to 2010 were collected from various sources. The model was fitted for each year individually, and we obtained model fitting R^2 ranging from 0.71 to 0.85, MPE from 1.73 to 2.50 $\mu\text{g m}^{-3}$, and RMSPE from 2.75 to 4.10 $\mu\text{g m}^{-3}$. In addition, we found cross validation R^2 ranging from 0.62 to 0.78, MPE from 2.00 to 3.01 $\mu\text{g m}^{-3}$, and RMSPE from 3.12 to 5.00 $\mu\text{g m}^{-3}$, indicating a good agreement between the estimated and observed values. Spatial trends show that high $\text{PM}_{2.5}$ levels occurred in urban areas and along major highways, while low concentrations appeared in rural or mountainous areas. A time series analysis was conducted to examine temporal trends of $\text{PM}_{2.5}$ concentrations in the study area from 2001 to 2010. The results showed that the $\text{PM}_{2.5}$ levels in the study area followed a generally declining trend from 2001 to 2010 and decreased about 20 % during the period. However, there was an exception of an increase in year 2005, which is attributed to elevated sulfate concentrations in the study area in

warm months of 2005. An investigation of the impact of wild and prescribed fires on $\text{PM}_{2.5}$ levels in 2007 suggests a positive relationship between them.

1 Introduction

Long-term exposure to $\text{PM}_{2.5}$ (particle size less than $2.5\mu\text{m}$ in the aerodynamic diameter) is associated with various adverse health outcomes including respiratory and cardiovascular diseases (Crouse et al., 2012; Peng et al., 2009). Due to the spatiotemporally continuous nature of fine particles, obtaining long-term and spatially-resolved distribution of $\text{PM}_{2.5}$ concentrations is important to reduce exposure misclassification and facilitate accurate epidemiological studies in the region. In addition, time series analyses of air pollution and human health have become the most common study design to compare day-to-day fluctuations of air pollution and corresponding fluctuations in health outcomes (Ito et al., 2007), requiring long-term $\text{PM}_{2.5}$ concentration estimates. Previous research examined temporal trends in $\text{PM}_{2.5}$ levels. For instance, Weber et al. (2003) investigated the temporal variations in $\text{PM}_{2.5}$ mass at the US Environmental Protection Agency (EPA) Atlanta Supersite Experiment in August 1999. So et al. (2007) examined long-term variation in $\text{PM}_{2.5}$ levels during two 12 month periods in Hong Kong. EPA (2011) reported temporal trends of annual and 24 h mean $\text{PM}_{2.5}$ concentrations at the national level from 2001 to 2010 and found that annual and 24 h mean $\text{PM}_{2.5}$ concentrations dropped 24 % and 28 %, respectively, within these ten years.

Another critical aspect is the investigation of spatial trends of $\text{PM}_{2.5}$ levels. Stationary ambient monitors have been established to measure ground-level $\text{PM}_{2.5}$ concentrations, yet those point measurements leave large areas uncovered, and therefore spatial variability is difficult to assess with those point measurements alone. In addition, these sites do not measure individual-specific exposure, and thus this approach inevitably introduce measurement errors that likely have substantial implications for interpreting epidemiological studies, especially for time-series analyses (Zeger et al., 2000).

25619

10 yr trend of $\text{PM}_{2.5}$ concentrations in the southeastern US

X. Hu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



On the other hand, aerosol observations from satellite remote sensing could substantially improve estimates of population exposure to $\text{PM}_{2.5}$ (van Donkelaar et al., 2010). As a result, satellite-retrieved aerosol optical depth (AOD) that measures light extinction by aerosols in the atmospheric column has been widely used to predict ground-

5 level $\text{PM}_{2.5}$ concentrations, considering its relatively low cost and large spatiotemporal coverage. A number of AOD products from sensors such as the Moderate Resolution Imaging Spectroradiometer (MODIS), the Multiangle Imaging SpectroRadiometer (MISR), and the Geostationary Operational Environmental Satellite Aerosol/Smoke Product (GASP) have been applied to $\text{PM}_{2.5}$ concentration prediction in previous studies

10 (Liu et al., 2007, 2009; Paciorek et al., 2008; Hu et al., 2013). Nevertheless, one of the limitations inherent with those AOD products is their coarse spatial resolutions. For instance, the spatial resolutions of AOD derived from MODIS and MISR are 10 km and 17.6 km, respectively. Although GASP has a spatial resolution of 4 km, the AOD retrievals are less precise than those from the polar-orbiting instruments due to a limited

15 information content (one spectral band) and relatively low signal-to-noise ratio of GOES sensor (Prados et al., 2007). Meanwhile, epidemiological studies typically have access to health data geo-coded to small geographical units (e.g., zip code and census block groups), many of which are substantially smaller than the spatial resolutions of MODIS and MISR. In addition, estimated $\text{PM}_{2.5}$ concentrations at coarse resolutions inevitably omit some details of spatial variability of $\text{PM}_{2.5}$ exposure and therefore have

20 fundamental limitations in the investigation of spatial trends of $\text{PM}_{2.5}$ levels. Hence, it is essential to use high resolution AOD retrievals to generate high spatial resolution $\text{PM}_{2.5}$ concentration estimates. Recently, a new AOD product retrieved by a multi-angle implementation of atmospheric correction (MAIAC) algorithm has been reported (Lyapustin et al., 2011b). MAIAC AOD has a spatial resolution of 1 km and thus has the ability to estimate $\text{PM}_{2.5}$ concentrations at that resolution. Moreover, the MAIAC AOD has been demonstrated to be strongly associated with monitored $\text{PM}_{2.5}$ levels in the

25 New England region (Chudnovsky et al., 2012).

10 yr trend of $\text{PM}_{2.5}$ concentrations in the southeastern US

X. Hu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures




Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Various statistical methods have been developed to establish the quantitative relationship between $PM_{2.5}$ and satellite-derived AOD including linear regression (Schafer et al., 2008; Wallace et al., 2007; Gupta and Christopher, 2009), many of which do not consider day-to-day variability. Lee et al. (2011) and Kloog et al. (2011) argued that the $PM_{2.5}$ -AOD relationship varies day-to-day, and the temporal variability needs to be accounted for in order to improve performance of the AOD-based prediction models. As a result, both studies developed a linear mixed effects model to incorporate daily calibration of the $PM_{2.5}$ -AOD relationship and obtained predictions with high accuracy. To move one step further, we introduce a geographically weighted regression (GWR) model as the second stage to account for the spatial variability in the $PM_{2.5}$ -AOD relationship.

The objective of this paper is first, to estimate spatiotemporally-resolved $PM_{2.5}$ concentrations in the study domain during the period between 2001 and 2010 using a two-stage model with MAIAC AOD as the primary predictor and meteorological and land use variables as the secondary predictors. Second, maps of annual mean $PM_{2.5}$ concentrations as well as the changes between 2001 and 2010 were generated from the daily estimates to visually illustrate spatial trends of annual $PM_{2.5}$ levels from 2001 to 2010. Third, time series analyses were conducted for the study domain and the Atlanta metro area using the monthly and annual mean $PM_{2.5}$ estimates to examine the 10 yr temporal trends of fine particle levels, and the underlying causes were discussed. Finally, potential impact of wild and prescribed fires on $PM_{2.5}$ levels in the south of our study region was investigated.

2 Materials and methods

2.1 Study area

The study area is approximately $600 \times 600 \text{ km}^2$ in the southeastern US, covering most of Georgia, Alabama, and Tennessee, and parts of North and South Carolina (Fig. 1).

The domain includes several large urban centers, numerous medium to small cities, and suburban and rural areas. Additionally, Zhang et al. (2010) reported that biomass burning contributes 13 % to PM_{2.5} mass annually in the southeastern US. Therefore, to investigate the impact of wild and prescribed fires on PM_{2.5} levels, two test sites of the same size (e.g., 18 × 18 km²) were selected in the south of the study area. Those two adjacent sites are both located in rural areas and away from any major emission source. One has an accumulated number of 472 fires from 2001 to 2010 (40 fires in 2007), and the other has a total number of 80 fires during the same period of time (8 fires in 2007).

2.2 PM_{2.5} measurements

The 24 h average PM_{2.5} concentrations from 2001 to 2010 collected from US Environmental Protection Agency (EPA) federal reference monitors (FRM) were downloaded from the EPA's Air Quality System Technology Transfer Network (<http://www.epa.gov/ttn/airs/airsaqs/>). PM_{2.5} concentrations less than 2 µg m⁻³ (~0.2–3 % of total data records) were discarded as they are below the established limit of detection (EPA, 2008a).

2.3 Remote sensing data

MAIAC retrieves aerosol parameters over land at 1 km resolution, which was accomplished by using the time series of MODIS measurements and simultaneous processing of a group of pixels in fixed 25 × 25 km² blocks (Lyapustin et al., 2011a, b, 2012). MAIAC uses a sliding window to collect up to 16 days of MODIS radiance observations over the same area and processes them to obtain surface parameters used for aerosol retrievals. To facilitate the time series analysis, MODIS data are initially gridded to a 1 km resolution in a selected projection. For this work, we used MODIS level 1B (calibrated and geometrically corrected) data from Collection 6 re-processing, which

removed major effects of temporal calibration degradation of Terra and Aqua, a necessary prerequisite for the trend analysis.

Validation showed that the MAIAC and operational Collection 5 MODIS Dark Target AOD have a similar accuracy over dark and vegetated surfaces, but also showed that MAIAC generally improves accuracy over brighter surfaces, including most urban areas (Lyapustin et al., 2011b). MAIAC AOD data from 2001 to 2010 were obtained from NASA Goddard Space Flight Center.

Zhang et al. (2012) found that Terra and Aqua may provide a good estimate of the daily average of AOD, thus the average of Aqua and Terra measurements can be used to predict $PM_{2.5}$ concentrations. In this study, Aqua (overpass at ~1:30 p.m. LT) and Terra (overpass at ~10:30 a.m. LT) MAIAC AOD values were first combined to improve spatial coverage. In a common MAIAC pixel, there might be only one MAIAC product from either Aqua or Terra, or both may be present. In the second case, when we combine them, the averaged value represents the mean of the AOD distribution from 10 a.m. to 2 p.m. LT, yet in the first case, AOD as an indicator of $PM_{2.5}$ abundance is biased towards the atmospheric condition either in the morning or early afternoon. To overcome this bias, Lee et al. (2011) defined a simple ratio between averaged Terra and Aqua AOD to estimate the missing AOD value. In this study, we fitted a linear regression to define the relationship between daily mean AOD values of MAIAC-Terra and MAIAC-Aqua. We used this regression to predict the missing AOD value (i.e., predict MAIAC-Terra AOD with the available MAIAC-Aqua AOD, and vice versa), then averaged the observed and the predicted AOD values together. Finally, we set an upper bound of 2.0 for the combined AOD to reduce potential cloud contamination (~0.05–0.1 % of total data records were excluded).

The fire detection data for the study region were obtained from the US Department of Agriculture (USDA) Forest Service's Remote Sensing Applications Center for 2001 through 2010 ([http://activefiremaps.fs.fed.us/gisdata.php?sensor=modis&extent=north america](http://activefiremaps.fs.fed.us/gisdata.php?sensor=modis&extent=north+america)).

10 yr trend of $PM_{2.5}$
concentrations in the
southeastern US

X. Hu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



These data are Terra and Aqua MODIS fire and thermal anomalies data from the official NASA MCD14ML product, Collection 5, Version 1.

2.4 Meteorological fields

The meteorological fields provided by the North American Land Data Assimilation System (NLDAS) Phase 2 were downloaded from the NLDAS website (<http://ldas.gsfc.nasa.gov/nldas/>). The spatial resolution of NLDAS meteorological data is 1/8th-degree (~13 km). Another meteorological dataset used in this study is the North American Regional Reanalysis (NARR). NARR is a long term, consistent, high-resolution climate dataset for North America (Mesinger et al., 2006). The spatial resolution of the NARR dataset is ~32 km. NLDAS provides most of the meteorological fields used in this analysis, including relative humidity, u-wind, and v-wind, while NARR provides another critical parameter: boundary layer height. To generate daytime meteorological fields corresponding to the MODIS overpass times, 3-hourly NARR measurements and hourly NLDAS measurements from 10 a.m. to 4 p.m. standard local time were averaged.

2.5 Land use variables

Elevation data were downloaded from the national elevation dataset (NED) (<http://ned.usgs.gov>). NED is the seamless elevation dataset covering the conterminous United States and is distributed by the US Geological Survey (USGS). The elevation data are downloaded at a spatial resolution of 1 arcsec (~30 m). The road data were obtained from ESRI StreetMap USA (Environmental Systems Research Institute, Inc., Redland, CA). The road data at level A1 (limited access highway) were extracted. Summed length of road segments were calculated for each 1 × 1 km² MAIAC grid cell, and grid cells with no roads were assigned zero. 2001 and 2006 Landsat-derived land cover maps covering the study area with a spatial resolution of 30 m were downloaded from the National Land Cover Database (NLCD) (<http://www.epa.gov/mrlc/nlcd->

2001.html). Forest cover maps were generated by assigning one to forest pixels and zero to others. Primary PM_{2.5} emissions (tons per year) were obtained from the 2002, 2005, and 2008 EPA National Emissions Inventory (NEI) facility emissions reports. Grid cells with multiple emission sources were assigned the summed value, and those with
 5 no emissions were assigned zero.

2.6 Data integration

All the data were first re-projected to the USA Contiguous Albers Equal Area Conic USGS coordinate system. For model fitting, a 1 × 1 km² square buffer was generated for each PM_{2.5} monitor site. Meteorological fields and AOD values were assigned to each
 10 PM_{2.5} monitor site using the nearest neighbor approach. Forest cover and elevation were averaged, while road length and point emissions were summed over the 1 × 1 km² square buffer. For PM_{2.5} prediction, the same procedure was performed for each 1 × 1 km² MAIAC grid cell.

2.7 Model structure

15 We developed a two-stage spatiotemporal model. The first stage is a linear mixed effects model with day-specific random intercepts and slopes for AOD, meteorological fields to account for the temporally varying relationship between PM_{2.5} and AOD (Eq. 1). The model structure can be expressed as

$$\begin{aligned} \text{PM}_{2.5,st} = & (b_0 + b_{0,t}) + (b_1 + b_{1,t})\text{AOD}_{st} + (b_2 + b_{2,t})\text{Meteorological Fields}_{st} \\ & + b_3\text{Elevation}_s + b_4\text{Major Roads}_s + b_5\text{Forest Cover}_s \\ & + b_6\text{Point Emissions}_s + \varepsilon_{st}(b_{0,t}, b_{1,t}, b_{2,t}) \sim N[(0, 0, 0), \Psi] \end{aligned} \quad (1)$$

20 where b_i and $b_{i,t}$ (day-specific) are the fixed and random intercept and slopes, respectively; PM_{2.5,st} is the measured ground level PM_{2.5} concentration (μg m⁻³) at site s in day t ; AOD_{st} is the MAIAC AOD value (unitless) at site s in day t ; Meteorologi-

25625

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



cal Fields_{st} is the meteorological parameters at site *s* in day *t* and may include Relative Humidity_{st}, Boundary Layer Height_{st}, Wind Speed_{st}, U-Wind_{st}, and V-Wind_{st}; Relative Humidity_{st} is the relative humidity (%) at site *s* in day *t*; Boundary Layer Height_{st} is the boundary layer height (m) at site *s* in day *t*; Wind Speed_{st} is the 2 m wind speed
 5 (ms⁻¹) at site *s* in day *t*; U-Wind_{st} is the east-west component of wind (ms⁻¹) at site *s* in day *t*; V-Wind_{st} is the north-south component of wind (ms⁻¹) at site *s* in day *t*; Elevation_s is elevation values (m) at site *s*; Major Roads_s is road length values (m) at site *s*; Forest Cover_s is forest cover values at site *s*; Point Emissions_s is point emissions (tyr⁻¹) at site *s*; and Ψ is an unstructured variance-covariance matrix for the random
 10 effects. The fixed effects affect the population mean and represent the average effects on PM_{2.5} concentration estimates for the entire period, while the random effects contribute to the covariance structure and account for the daily variability in associations between dependent and independent variables.

The second stage is a geographically weighted regression (GWR) model that can
 15 generate a continuous surface of estimates for each parameter at each location instead of a universal value for all observations. We fitted a monthly GWR model to calibrate the spatial variability within the PM_{2.5}-AOD relationship, and the model can be expressed as

$$\text{PM}_{2.5_resi_{st}} = \beta_{0,s} + \beta_{1,s}\text{AOD}_{st} + \varepsilon_{st} \quad (2)$$

20 where PM_{2.5_resi_{st}} denotes the residuals from the stage one model at site *s* in month *t*, AOD_{st} is the MAIAC AOD value (unitless) at site *s* in month *t*, and $\beta_{0,s}$ and $\beta_{1,s}$ denote the location-specific intercept and slope, respectively.

It should be noted that the model was fitted for each year individually, and therefore the predictors used in the model may vary for different years. The model structures
 25 were determined by comparison of performance of models using different predictors to ensure that relatively better prediction accuracy can be achieved. To assess the goodness of fit of the model, various statistical indicators such as the coefficient of determination (R^2), mean prediction error (MPE), and square root of the mean squared

10 yr trend of PM_{2.5} concentrations in the southeastern US

X. Hu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



prediction errors (RMSPE) were calculated between the fitted $\text{PM}_{2.5}$ concentrations from the model and the observations. In addition, a 10-fold cross validation (CV) technique was adopted to assess the potential model over-fitting. That is, the model could perform better on the data used to fit the model than unobserved data. The entire model-fitting dataset was randomly split into ten subsets with approximately 10 % of the total data records in each subset. In each round of cross validation, we selected one subset (10 % of the data) as testing samples and used the remaining nine subsets (90 % of the data) to fit the model. Predictions of the held-out subset (10 % of the data) were made from the fitted model. The process was repeated ten times until every subset was tested. Statistical indicators such as R^2 , MPE, and RMSPE were calculated between the CV predicted concentrations and the observations. The model over-fitting assessment was conducted by the comparison between CV and model-fitting statistics. A relative accuracy value was also calculated for each year to make validation results comparable among different years.

The daily $\text{PM}_{2.5}$ concentrations were estimated using the model for each year individually. The maps of annual mean $\text{PM}_{2.5}$ concentrations as well as the changes between 2001 and 2010 for the study domain and the Atlanta metro area were generated using the daily estimates to visually examine spatial trends of $\text{PM}_{2.5}$ levels from 2001 to 2010. Moreover, time series analyses were conducted by year and month, respectively to quantitatively investigate the 10 yr temporal trends of fine particle levels in the study domain and the Atlanta metro area. Finally, an examination of the impact of wild and prescribed fires on $\text{PM}_{2.5}$ levels was also conducted on two test sites.

3 Results

3.1 Descriptive statistics

The descriptive statistics of variables used in fitting the models are listed in Table 1. The annual mean $\text{PM}_{2.5}$ concentrations ranged from 11.03 to $15.63 \mu\text{m}^{-3}$ between

2001 and 2010, the highest occurred in 2005, and the lowest appeared in 2009. The annual mean AOD values ranged from 0.20 to 0.28 during the same period of time. Table 1 also shows that land use variables and meteorological fields vary from year to year within the data.

3.2 Results of model-fitting and validation

The model-fitting and CV statistics (e.g. R^2 , MPE, and RMSPE) are listed in Table 2. The results show that R^2 ranges from 0.71 to 0.85, MPE is from 1.73 to 2.50 $\mu\text{g m}^{-3}$, RMSPE ranges from 2.75 to 4.10 $\mu\text{g m}^{-3}$, and relative accuracy ranges from 72.9 % to 80.7 %, which indicate a good fit between predicted values from the fitted models and the observations. In addition, CV statistics results suggest that model over-fitting is present; that is, R^2 decreases, while MPE and RMSPE increase from model fitting to cross validation, yet the differences are relatively small for all the years. For instance, R^2 and relative accuracy have an average decrease of 0.08 and 4.21 %, respectively, while MPE and RMSPE have an average increase of 0.39 and 0.60 $\mu\text{g m}^{-3}$, respectively through all the years. Moreover, a regression with zero intercept (Fig. 2) was performed to fit the predicted values against the observations. The figure shows that at high concentration levels, both model fitting and cross validation under-predicted the $\text{PM}_{2.5}$ concentrations by 3–7 % (e.g. fitted/CV $\text{PM}_{2.5}$ = 97 % to 93 % observed $\text{PM}_{2.5}$).

3.3 Spatial trends of $\text{PM}_{2.5}$ concentrations

Figure 3 illustrates the $\text{PM}_{2.5}$ concentration estimates at 1 km spatial resolution in the study area. The annual mean estimated concentrations are 13.97, 13.90, 13.35, 13.31, 15.19, 13.73, 13.22, 11.34, 10.58, and 11.22 $\mu\text{g m}^{-3}$ for year 2001 through 2010, respectively. The patterns of $\text{PM}_{2.5}$ distribution are very similar for all the years. High concentrations appear in large urban centers and along major highways, while low concentrations occur in rural and mountainous areas. In addition, high $\text{PM}_{2.5}$ levels also occur in the south of the study domain. This area is primarily occupied by agriculture

land, and high agricultural emissions may lead to high concentrations of fine particles. As reported by previous studies, ammonia (NH₃) and nitrogen oxides (NO_x) generated by agricultural activities, such as farm vehicles, domestic and farm animals, and fertilizer applications, can significantly increase the number of suspended particles (Kurvits and Marta, 1998). In addition, biomass burning also contributes to emissions of fine particles in the region (Zhang et al., 2010). Figure 4 shows that the pattern of ground PM_{2.5} measurements from FRM monitors corresponds well with that of our estimated concentrations.

To take advantage of the high spatial resolution of the MAIAC data, a map of PM_{2.5} estimates in Atlanta metro area was also generated for each year (Fig. 5). The annual mean estimates from 2001 to 2010 are 15.10, 14.64, 14.00, 14.54, 15.63, 14.39, 14.14, 11.78, 10.98, and 11.65 µg m⁻³, respectively. Those maps distinctly show that high PM_{2.5} levels occur in areas with high urban land use and along major highways, while low concentrations appear in forest and recreational areas, suggesting an underlying positive relationship between air pollution levels and urban development.

Figure 6 shows the percent changes in PM_{2.5} concentrations from 2001 to 2010. Figure 6a illustrates the spatial trend of changes in PM_{2.5} levels in the study region. The results show that PM_{2.5} levels in most of the areas decreased from 0 to 25 %, and large parts of the areas had decreases exceeding 25 % and up to 50 %. Larger decreases occurred in areas with generally higher pollution levels such as Atlanta metro area and along major highways, which might be due to recently enacted emission reduction program (EPA, 2011), since majority of the emission sources are located in or near urban areas and along major highways. Mitigation of fine particles has been effected through control of direct PM_{2.5} emissions from both stationary and mobile sources (e.g., through installation of scrubbers and filters and the use of alternative fuels and electric vehicles) (EPA, 2007). One exception is the mountainous area in the northeast of our domain with generally low PM_{2.5} levels. The decreases of PM_{2.5} concentrations in this region from 2001 to 2010 were also phenomenal, PM_{2.5} levels decreased from 25 % to 50 % in most of the region, and some areas had decreases exceeding 50 %.

10 yr trend of PM_{2.5} concentrations in the southeastern US

X. Hu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



By checking 2002 and 2008 EPA NEI facility emissions reports, this is probably due to dramatically reduced number of emission sources in the region during the period. Figure 6b illustrates the percent changes in $PM_{2.5}$ levels in the Atlanta Metro area. Once again, the spatial trend shows that larger decreases (25 % to 50 %) primarily occurred in urban built-up areas and along major highways, while small decreases (0 to 25 %) appeared in forest or recreational areas with generally lower pollution levels. It is reasonable because the increase or decrease of emissions from emission sources might be the major driver causing fluctuations of $PM_{2.5}$ levels in this region, while most of those sources are unlikely to be located in forest and recreational areas. Two pixels with large changes (increasing more than 25 % and decreasing more than 50 %) were identified (in blue and red circles). The large decrease of $PM_{2.5}$ concentration in the blue pixel was because of the large reduction of particle emissions from power plants located within that pixel during the period between 2001 and 2010. Likewise, the large increase of $PM_{2.5}$ concentration in the red pixel was due to the addition of one emission source that did not exist in year 2001.

3.4 Temporal trends of $PM_{2.5}$ concentrations

A time series analysis was conducted to quantitatively examine temporal trends of $PM_{2.5}$ levels in the study area as well as the Atlanta metro area during the period between 2001 and 2010 (Fig. 7). The results show that the estimated $PM_{2.5}$ concentrations are smaller than the monitored observations, and underestimation by our model has occurred, especially in areas with high pollution levels (e.g., the Atlanta metro area), which agrees with our model validation results. The average underestimation is $0.99 \mu g m^{-3}$ for the study domain and $1.82 \mu g m^{-3}$ for the Atlanta metro area. The $PM_{2.5}$ levels in the study region as well as the Atlanta metro area followed a generally declining trend, especially after year 2005. From 2001 to 2010, the annual mean $PM_{2.5}$ concentration decreased about 20 % in the study area and 23 % in the Atlanta metro area, which is in line with the findings documented in the US EPA report on particle pollution (EPA, 2011). Both EPA's and our results illustrate a peak of $PM_{2.5}$ levels in

25630

ACPD

13, 25617–25648, 2013

10 yr trend of $PM_{2.5}$ concentrations in the southeastern US

X. Hu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



year 2005, and this phenomenon might be attributed to the increase of sulfate concentrations emitted from electric utilities and industrial boilers during the warm months (e.g, from May to September) of 2005 (EPA, 2008b). In addition, the steady decrease of PM_{2.5} levels after year 2005 is due to the emissions reduction programs that have been enacted recently (EPA, 2011, 2007).

3.5 The impact of wild and prescribed fires on PM_{2.5} concentrations

Zeng et al. (2008) suggested that prescribed fire emissions can result in a daily increase of PM_{2.5} mass up to 25 µg m⁻³, indicating high impact of fires on PM_{2.5} levels. Figure 8 illustrates the seasonal distribution of the number of fires in the study region, which shows that most wild and prescribed fires occurred in spring and fall, and the highest number of fires appeared in spring 2007. To assess the impact of fires on PM_{2.5} levels, we selected year 2007 to conduct a case study as fires in the southeastern US in 2007 has been examined by previous research using ground monitoring data (Zhang et al., 2010). The differences of number of fires and annual mean PM_{2.5} concentrations between two test sites were calculated and illustrated in Fig. 9. It shows that in most of the cases, the peaks and valleys of the differences of number of fires correspond well with the peaks and valleys of the differences of PM_{2.5} concentrations (indicated by black arrows), indicating an underlying positive relationship between fires and PM_{2.5} concentrations and suggesting that the fires may have impact on fine particle levels in the south of our study region. However, exceptions exist. For example, in April 2007, a day in which a large difference of number of fires occurred did not show a higher difference of PM_{2.5} concentrations than its neighboring days. In addition, in many days in which there are no differences of number of fires, the differences of PM_{2.5} concentrations between two test sites still exist. Such situations might be attributed to several factors: first, due to the lack of data, our analysis only used the number of fires without fire scale and intensity information and thus might be biased. Second, although the contributions of fires to PM_{2.5} levels are quite significant (~13 % annually) and vary by seasons in the southeastern US, fires might not be major drivers and there are other,

25631



often more substantial, contributors to PM_{2.5} concentrations in the region (Zhang et al., 2010).

4 Discussion

A major strength of this study is that we used 1 km spatial resolution PM_{2.5} estimates derived from 1 km resolution MAIAC AOD to investigate spatiotemporal trends of PM_{2.5} concentrations in the study area as well as in the Atlanta metro area. PM_{2.5} estimates at finer resolutions are more suitable for investigation of spatial trends than those at coarser resolutions derived from other AOD products (e.g., MODIS and MISR), because estimates at coarser scales inevitably omit local spatial details. Our results are capable of showing PM_{2.5} concentrations and changes within a 1 × 1 km² area, which are very useful for air pollution studies at local scales. For instance, spatial trends of changes in PM_{2.5} concentrations in the Atlanta metro area show distinct patterns of higher PM_{2.5} reduction in areas with generally higher pollution levels (e.g., urban built-up areas and along major highways). Some of the changes may be directly associated with the occurrence or disappearance of one or more emission sources as well as the increase or decrease of emissions from those sources. Although high resolution PM_{2.5} estimates can provide more details to examine spatial trends, difficulties lie in validation to ground monitoring. More ground measurements at specific locations are needed to further validate the results.

Our results of temporal trends of PM_{2.5} concentrations correspond well with EPA's results (EPA, 2011). However, results also show that our predicted PM_{2.5} concentrations underestimated the ground measurements for almost all the years, which is expected. Most of the EPA FRM monitors are located in or near urban areas with generally high PM_{2.5} levels. On the other hand, the temporal trends of PM_{2.5} concentrations estimated from satellite AOD include the entire study domain and account for both urban and rural estimates, and therefore might more thoroughly represent the true fluctuations of

Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper

ACPD

13, 25617–25648, 2013

10 yr trend of PM_{2.5} concentrations in the southeastern US

X. Hu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



regional fine particle levels, and further research will continue to explore these associations.

In addition, long-term $\text{PM}_{2.5}$ estimates at high spatial resolutions might be better suited for incorporating fire data in order to examine the impact of fires on fine particle levels, since fires might occur far away from ground monitors. Our results show that fires, to some extent, contribute to $\text{PM}_{2.5}$ concentrations in our study region. However, they might not be the major contributors to fine particle levels. It should be noted that our results are preliminary and might contain a certain degree of bias, since no fire scale and intensity data were incorporated in the analysis. However, in this paper, we primarily attempt to examine the 10 yr spatial and temporal trends of fine particle levels in the region using high spatial resolution $\text{PM}_{2.5}$ estimates as well as the possible causes for high $\text{PM}_{2.5}$ concentrations in the south of our domain. Hence, quantification of the contribution of fires to $\text{PM}_{2.5}$ levels was beyond the scope of this analysis.

5 Conclusions

In this paper, we used a two-stage spatiotemporal model incorporating MAIAC AOD data, meteorological fields, and land use variables to estimate $\text{PM}_{2.5}$ concentrations at 1 km spatial resolution and investigated the 10 yr spatial and temporal trends of $\text{PM}_{2.5}$ levels in our study region. The results show a reasonable spatial pattern of $\text{PM}_{2.5}$ levels in the study area as well as in the Atlanta metro area. For instance, high concentrations occur in large urban centers and along major highways, while low concentrations appear in rural and mountainous area. $\text{PM}_{2.5}$ estimates at high spatial resolutions can provide more details in small geographic regions and reduce exposure misclassification in air pollution and epidemiological studies. The spatial trends of changes in $\text{PM}_{2.5}$ concentrations indicate that higher pollution reduction occurred in areas with generally higher $\text{PM}_{2.5}$ levels (e.g. in urban areas and along major highways), while areas with generally lower pollution levels (e.g., in forest and recreational areas) had lower and moderate reduction of fine particle concentrations.

25633

ACPD

13, 25617–25648, 2013

10 yr trend of $\text{PM}_{2.5}$ concentrations in the southeastern US

X. Hu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Our time series analysis results indicate that the PM_{2.5} levels decreased about 20 % in the study region and 23 % in the Atlanta Metro area during the period between 2001 and 2010, especially after year 2005. In addition, an analysis of the impact of wild and prescribed fires on PM_{2.5} concentrations shows that the peaks and valleys of the differences of the number of fires corresponds well with the peaks and valleys of the differences of PM_{2.5} levels between two test sites, suggesting a positive relationship between fires and fine particle exposure that merits further investigation.

Acknowledgements. This work was partially supported by NASA Applied Sciences Program (grant no. NNX09AT52G and NNX11AI53G). In addition, this publication was made possible by USEPA grant R834799. Its contents are solely the responsibility of the grantee and do not necessarily represent the official views of the USEPA. Further, USEPA does not endorse the purchase of any commercial products or services mentioned in the publication.

References

- Chudnovsky, A. A., Kostinski, A., Lyapustin, A., and Koutrakis, P.: Spatial scales of pollution from variable resolution satellite imaging, *Environ. Pollut.*, 172, 131–138, doi:10.1016/j.envpol.2012.08.016, 2012.
- Crouse, D. L., Peters, P. A., van Donkelaar, A., Goldberg, M. S., Villeneuve, P. J., Brion, O., Khan, S., Atari, D. O., Jerrett, M., Pope, C. A., Brauer, M., Brook, J. R., Martin, R. V., Stieb, D., and Burnett, R. T.: Risk of non accidental and cardiovascular mortality in relation to long-term exposure to low concentrations of fine particulate matter: a Canadian National-Level Cohort study, *Environ. Health Persp.*, 120, 708–714, doi:10.1289/ehp.1104049, 2012.
- Gupta, P. and Christopher, S. A.: Particulate matter air quality assessment using integrated surface, satellite, and meteorological products: Multiple regression approach, *J. Geophys. Res.-Atmos.*, 114, D14205, doi:10.1029/2008jd011496, 2009.
- Hu, X., Waller, L. A., Al-Hamdan, M. Z., Crosson, W. L., Estes Jr, M. G., Estes, S. M., Quattrochi, D. A., Sarnat, J. A., and Liu, Y.: Estimating ground-level PM_{2.5} concentrations in the southeastern U.S. using geographically weighted regression, *Environ. Res.*, 121, 1–10, doi:10.1016/j.envres.2012.11.003, 2013.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Ito, K., Thurston, G. D., and Silverman, R. A.: Characterization of PM_{2.5}, gaseous pollutants, and meteorological interactions in the context of time-series health effects models, *J. Expo. Sci. Env. Epid.*, 17, S45–S60, 2007.
- Kurvits, T. and Marta, T.: Agricultural NH₃ and NO_x emissions in Canada, *Environ. Pollut.*, 102, 187–194, doi:10.1016/S0269-7491(98)80032-8, 1998.
- 5 Lee, H. J., Liu, Y., Coull, B. A., Schwartz, J., and Koutrakis, P.: A novel calibration approach of MODIS AOD data to predict PM_{2.5} concentrations, *Atmos. Chem. Phys.*, 11, 7991–8002, doi:10.5194/acp-11-7991-2011, 2011.
- Liu, Y., Franklin, M., Kahn, R., and Koutrakis, P.: Using aerosol optical thickness to predict ground-level PM_{2.5} concentrations in the St. Louis area: a comparison between MISR and MODIS, *Remote Sens. Environ.*, 107, 33–44, doi:10.1016/j.rse.2006.05.022, 2007.
- 10 Liu, Y., Paciorek, C. J., and Koutrakis, P.: Estimating regional spatial and temporal variability of PM_{2.5} concentrations using satellite data, meteorology, and land use information, *Environ. Health Persp.*, 117, 886–892, 2009.
- 15 Lyapustin, A., Martonchik, J., Wang, Y. J., Laszlo, I., and Korkin, S.: Multiangle implementation of atmospheric correction (MAIAC): 1. Radiative transfer basis and look-up tables, *J. Geophys. Res.-Atmos.*, 116, D03210, doi:10.1029/2010jd014985, 2011a.
- Lyapustin, A., Wang, Y., Laszlo, I., Kahn, R., Korkin, S., Remer, L., Levy, R., and Reid, J. S.: Multiangle implementation of atmospheric correction (MAIAC): 2. Aerosol algorithm, *J. Geophys. Res.-Atmos.*, 116, D03211, doi:10.1029/2010jd014986, 2011b.
- 20 Lyapustin, A. I., Wang, Y., Laszlo, I., Hilker, T., Hall, F. G., Sellers, P. J., Tucker, C. J., and Korkin, S. V.: Multi-angle implementation of atmospheric correction for MODIS (MAIAC): 3. Atmospheric correction, *Remote Sens. Environ.*, 127, 385–393, doi:10.1016/j.rse.2012.09.002, 2012.
- 25 Mesinger, F., DiMego, G., Kalnay, E., Mitchell, K., Shafran, P. C., Ebisuzaki, W., Jovic, D., Woollen, J., Rogers, E., and Berbery, E. H.: North American regional reanalysis, *B. Am. Meteorol. Soc.*, 87, 343–360, 2006.
- Paciorek, C. J., Liu, Y., Moreno-Macias, H., and Kondragunta, S.: Spatiotemporal associations between GOES aerosol optical depth retrievals and ground-level PM_{2.5}, *Environ. Sci. Technol.*, 42, 5800–5806, doi:10.1021/es703181j, 2008.
- 30 Peng, R. D., Bell, M. L., Geyh, A. S., McDermott, A., Zeger, S. L., Samet, J. M., and Dominici, F.: Emergency admissions for cardiovascular and respiratory diseases and the

10 yr trend of PM_{2.5} concentrations in the southeastern US

X. Hu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- chemical composition of fine particle air pollution, *Environ. Health Persp.*, 117, 957–963, doi:10.1289/ehp.0800185, 2009.
- Prados, A. I., Kondragunta, S., Ciren, P., and Knapp, K. R.: GOES Aerosol/Smoke Product (GASP) over North America: comparisons to AERONET and MODIS observations, *J. Geophys. Res.*, 112, D15201, doi:10.1029/2006jd007968, 2007.
- 5 Schafer, K., Harbusch, A., Emeis, S., Koepke, P., and Wiegner, M.: Correlation of aerosol mass near the ground with aerosol optical depth during two seasons in Munich, *Atmos. Environ.*, 42, 4036–4046, doi:10.1016/j.atmosenv.2008.01.060, 2008.
- So, K. L., Guo, H., and Li, Y. S.: Long-term variation of PM_{2.5} levels and composition at rural, urban, and roadside sites in Hong Kong: increasing impact of regional air pollution, *Atmos. Environ.*, 41, 9427–9434, doi:10.1016/j.atmosenv.2007.08.053, 2007.
- 10 van Donkelaar, A., Martin, R. V., Brauer, M., Kahn, R., Levy, R., Verduzco, C., and Villeneuve, P. J.: Global estimates of ambient fine particulate matter concentrations from satellite-based aerosol optical depth: development and application, *Environ. Health Persp.*, 118, 847–855, doi:10.1289/ehp.0901623, 2010.
- Wallace, J., Kanaroglou, P., and Ieee: An investigation of air pollution in southern Ontario, Canada, with MODIS and MISR aerosol data, in: *Igarss: 2007 Ieee International Geoscience and Remote Sensing Symposium*, Vol. 1–12 – Sensing and Understanding Our Planet, IEEE International Symposium on Geoscience and Remote Sensing (IGARSS), Ieee, New York, July 23–28, 4311–4314, 2007.
- 20 Weber, R., Orsini, D., Duan, Y., Baumann, K., Kiang, C. S., Chameides, W., Lee, Y. N., Brechtel, F., Klotz, P., Jongejan, P., ten Brink, H., Slanina, J., Boring, C. B., Genfa, Z., Dasgupta, P., Hering, S., Stolzenburg, M., Dutcher, D. D., Edgerton, E., Hartsell, B., Solomon, P., and Tanner, R.: Intercomparison of near real time monitors of PM_{2.5} nitrate and sulfate at the US Environmental Protection Agency Atlanta Supersite, *J. Geophys. Res.-Atmos.*, 108, 8421, doi:10.1029/2001jd001220, 2003.
- 25 Zeger, S. L., Thomas, D., Dominici, F., Samet, J. M., Schwartz, J., Dockery, D., and Cohen, A.: Exposure measurement error in time-series studies of air pollution: concepts and consequences, *Environ. Health Persp.*, 108, 419–426, doi:10.2307/3454382, 2000.
- 30 Zeng, T., Wang, Y., Yoshida, Y., Tian, D., Russell, A. G., and Barnard, W. R.: Impacts of prescribed fires on air quality over the southeastern United States in spring based on modeling and ground/satellite measurements, *Environ. Sci. Technol.*, 42, 8401–8406, doi:10.1021/es800363d, 2008.

ACPD

13, 25617–25648, 2013

10 yr trend of PM_{2.5} concentrations in the southeastern US

X. Hu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Zhang, X., Hecobian, A., Zheng, M., Frank, N. H., and Weber, R. J.: Biomass burning impact on PM_{2.5} over the southeastern US during 2007: integrating chemically speciated FRM filter measurements, MODIS fire counts and PMF analysis, Atmos. Chem. Phys., 10, 6839–6853, doi:10.5194/acp-10-6839-2010, 2010.
- ⁵ Zhang, Y., Yu, H., Eck, T. F., Smirnov, A., Chin, M., Remer, L. A., Bian, H., Tan, Q., Levy, R., Holben, B. N., and Piazzolla, S.: Aerosol daytime variations over North and South America derived from multiyear AERONET measurements, J. Geophys. Res.-Atmos., 117, D05211, doi:10.1029/2011jd017242, 2012.

Table 1. Descriptive statistics (2001–2010).

| Var. | Min | Std. dev. | Max | Mean |
|---|-----------------|---------------|-----------------|----------------|
| PM _{2.5} (µg m ⁻³) | 2.0–2.6 | 5.31–8.64 | 50.1–145.0 | 11.03–15.63 |
| Boundary layer height (m) | 215–464 | 347–493 | 2605–3405 | 1146–1464 |
| Relative humidity (%) | 13.9–26.2 | 8.7–11.3 | 86.8–93.1 | 46.8–59.9 |
| U-wind (m s ⁻¹) | –9.44 to –6.30 | 2.62–3.20 | 10.22–16.85 | 0.82–1.47 |
| V-wind (m s ⁻¹) | –12.60 to –9.34 | 2.62–3.00 | 8.45–11.84 | –0.74 to –0.09 |
| Wind speed (m s ⁻¹) | 0.03–0.12 | 1.81–2.13 | 12.76–18.06 | 3.48–3.99 |
| Forest cover 2001 | 0 | 0.16–0.18 | 0.83 | 0.14–0.17 |
| Forest cover 2006 | 0 | 0.15–0.17 | 0.79 | 0.14–0.16 |
| Road length (m) | 0 | 187.29–230.81 | 1012.97–1078.09 | 58.05–82.92 |
| Elevation (m) | 46.78 | 126.82–141.65 | 811.63–822.82 | 227.74–249.10 |
| Point emissions 2002 (tyr ⁻¹) | 0 | 56.64–70.39 | 364.42 | 11.13–16.46 |
| Point emissions 2005 (tyr ⁻¹) | 0 | 150.89–188.63 | 985.48 | 26.90–40.84 |
| Point emissions 2008 (tyr ⁻¹) | 0 | 15.89–19.72 | 101.74 | 3.14–4.54 |
| AOD | 0–0.01 | 0.16–0.26 | 1.42–1.96 | 0.20–0.28 |

10 yr trend of PM_{2.5} concentrations in the southeastern US

X. Hu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 2. Model validation.

| Year | Model fitting | | | | Cross validation | | | |
|------|---------------|---------------------------------|-----------------------------------|---------------------------------------|------------------|---------------------------------|-----------------------------------|---------------------------------------|
| | R^2 | MPE ($\mu\text{g m}^{-3}$) | RMSPE ($\mu\text{g m}^{-3}$) | Relative accuracy (%) ^a | R^2 | MPE ($\mu\text{g m}^{-3}$) | RMSPE ($\mu\text{g m}^{-3}$) | Relative accuracy (%) ^a |
| 2001 | 0.78 | 2.50 | 4.10 | 72.9 | 0.67 | 3.01 | 5.00 | 67.0 |
| 2002 | 0.84 | 2.10 | 2.98 | 80.7 | 0.75 | 2.62 | 3.75 | 75.7 |
| 2003 | 0.85 | 1.95 | 2.77 | 80.4 | 0.76 | 2.42 | 3.47 | 75.4 |
| 2004 | 0.85 | 1.97 | 2.77 | 80.3 | 0.77 | 2.40 | 3.37 | 76.1 |
| 2005 | 0.84 | 2.23 | 3.17 | 79.7 | 0.78 | 2.64 | 3.76 | 75.9 |
| 2006 | 0.85 | 2.02 | 2.90 | 80.6 | 0.78 | 2.43 | 3.49 | 76.6 |
| 2007 | 0.79 | 2.26 | 3.75 | 74.0 | 0.71 | 2.64 | 4.39 | 69.6 |
| 2008 | 0.74 | 1.93 | 3.13 | 75.4 | 0.67 | 2.21 | 3.53 | 72.3 |
| 2009 | 0.71 | 1.73 | 2.88 | 73.9 | 0.62 | 2.00 | 3.28 | 70.3 |
| 2010 | 0.73 | 1.90 | 2.75 | 77.6 | 0.66 | 2.15 | 3.12 | 74.5 |

^a Relative accuracy is defined as $100\% - \text{RMSPE}/\text{the mean PM}_{2.5} \text{ concentration}$.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 1. Study area and two test sites.

10 yr trend of PM_{2.5} concentrations in the southeastern US

X. Hu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 2. Model validation. **(a)** Model fitting; **(b)** cross validation.

10 yr trend of PM_{2.5} concentrations in the southeastern US

X. Hu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 3. Annual mean PM_{2.5} concentration predictions in the study area.

10 yr trend of PM_{2.5} concentrations in the southeastern US

X. Hu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 4. Annual mean $\text{PM}_{2.5}$ concentration measured from ground FRM monitors.

10 yr trend of $\text{PM}_{2.5}$ concentrations in the southeastern US

X. Hu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 5. Annual mean $PM_{2.5}$ concentration predictions in the Atlanta metro area.




Fig. 6. The percent change of $\text{PM}_{2.5}$ concentrations in the study area **(a)** and the Atlanta metro area **(b)** from 2001 to 2010.




Fig. 7. Time series analysis of annual and monthly $\text{PM}_{2.5}$ concentrations. **(a)** Study area; **(b)** Atlanta metro area.




Fig. 8. Time series analysis of number of fires that occurred in the study area from 2001 to 2010.




Fig. 9. Time series analysis of relationship between differences of number of fires and differences of $\text{PM}_{2.5}$ concentrations for year 2007 (Only months with differences of number of fires were plotted, and the coincidences were indicated by black arrows).